

Late-Stage Prostate Cancer Incidence, by Year, Maine, 2004-2009

Year	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
2004	174	27.3	23.4 - 31.6	24.1	20.6 - 28.1
2005	167	26.1	22.3 - 30.4	23.3	19.9 - 27.2
2006	200	31.1	27.0 - 35.8	27.4	23.6 - 31.5
2007	185	28.8	24.8 - 33.2	24.1	20.7 - 27.9
2008	210	32.6	28.3 - 37.3	26.3	22.8 - 30.3
2009	160	24.9	21.2 - 29.0	19.6	16.6 - 23.0

Late-Stage Prostate Cancer Incidence, by 3-year averages, Maine, 2004-2009

Year	Count	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
2004-2006	541	28.2	25.8 - 30.6	25.0	22.9 - 27.2
2005-2007	552	28.7	26.3 - 31.2	24.9	22.9 - 27.2
2006-2008	595	30.8	28.4 - 33.4	25.9	23.8 - 28.1
2007-2009	557	28.8	26.5 - 31.3	23.4	21.4 - 25.5

Data Source: Maine Cancer Registry, November 2011 NPCR data submission.

Count: Number of cases diagnosed with malignant cancer during that time period.

Late-Stage: Cancers with 'regional' or 'distant' spread according to 2000 Derived Summary Stage criteria.

Prostate Cancer: SEER cancer site recodes: 28010 (ICD-O-3 codes: C619).

Crude rates are new cases per 100,000 population.

95% CI: 95% confidence interval of the rate.

Rates are calculated using SEER*Stat Version 7.0.5.